樹豆品種 (系) 雜交選育

陳振義

行政院農委會臺東區農業改良場

摘　要

樹豆在臺東原住民部落地區普遍種植，為原住民的傳統作物，亦被視為一種健康食品。目前國內地方品系均有成熟期不一致且低產之現象，造成採收及管理上的困擾，為解決上述之問題，擬以人工雜交方式雜交授粉，並於後代選育出高產及成熟期一致之雜交新品種，供農民種植。本場 101 年以人工雜交方式獲得其雜交種子，成功率為 23.6%，期望將來於雜交後代選育出高產及成熟期一致之新品種，供農民種植以增加農民收益。

關鍵詞：樹豆、雜交、選育、去雄、授粉。

前　言

樹豆(*Cajanus cajan* (L.) Millsp.) 為亞洲、非洲、拉丁美洲及加勒比海地區等熱帶發展中國家重要蛋白質補充來源。其乾豆產量占世界豆類產量排名第六，未成熟的種子可當作新鮮或加工用的蔬菜，成熟的種子可磨成粉，枝條亦能當作綠肥、牧草、覆蓋作物，尤其適合貧瘠及乾旱地區生產。在諸多的利用價值及極少的成本之下，樹豆在乾旱或半乾旱農業環境地區，已成為理想的作物。

樹豆以印度栽培面積最大，超過 70%以上，並擁有大多數的遺傳物質及特性，極可能為樹豆的起源地；其他重要的國家包括孟買、烏干達、肯亞、多明尼加、巴拿馬、波多黎各及西印度群島等地區。臺東地區之樹豆則主要為原住民傳統作物之一。

樹豆屬於常異交作物，其花序屬於無限型生長，以致樹豆開花、結莢及豆莢成熟時間參差不一，即使同一株也要相差 15-30 天，在採收上無法一致。為解決上述之問題，擬以人工雜交方式，於樹豆開花期進行去雄、雜交授粉，並於雜交之後代選育出高產及成熟期一致之新品種，供農民種植以增加農民收益。

材料與方法

1.試驗材料：以本場選育出優良地方品種(系)：臺東 1 號(白色)、TS-221(紅色)、 臺東 2 號(茶褐色)、臺東 3 號(黑色)及 TS-303(紅色)等 5 品種(系)為親本，作為雜交的材料。

2.親本組合：分別以臺東 1 號×TS-221、臺東 2 號×TS-221 及臺東 3 號×TS303 等三個組合進行雜交，共計 6 個組合。

3.親本種植：將上述各品種(系)於 101 年 4 月 16 日定植於本場豐里試驗地樹豆雜交圃，進行父、母本田間管理及開花前搭設簡易網室。

4.去雄授粉：於開花期間進行父、母本去雄、授粉工作，並調查各親本雜交成功率，記錄雜交後幼莢生長情形及記錄授粉當日溫度，以供日後之參考。

結果與討論

乾燥晴朗的氣候有利樹豆開花，陰天潮濕則會減少其開花數目，均勻且持續的光照是授粉及莢果形成所不可缺少的條件。樹豆在巴基斯坦，三月份最高溫度 38-40℃的條件下仍可授粉成功，但是到四月溫度超過 40℃

就會影響其授粉。樹豆的花開放的順序是由一個分支的基部開始向上開放。大小不同的花序，在不同類型每個花序、花朵可能有多達 10 個。通常在同一花序，每次有兩朵花同時開，直到莢果成熟。每一花序可以結 2 至 8 個莢果，莢果數量在分枝上至頂端上會逐漸減少。若於低節點先前的花序所開放的花和莢果都脫落，則節點上的莢果會再繼續結莢，以彌補脫落的損失。這種補償機制有助於延長頂端分生組織生長及開花期。

本場自 99-100 年進行樹豆品種(系)雜交選育工作，育種材料由本場選育之優良地方品系臺東 1 號(白色)、S-14(白色)、臺東 2 號(茶褐色)、S-76(白色)、臺東 3 號(黑色)及 S-95(白色)等 6 個優良地方品種(系)為親本，作為雜交的材料。以臺東 1 號×S-14、臺東 2 號×S-76 及臺東 3 號×S-95 等三個組合進行雜交，共計 6 個組合。101 年以混合選種法播種後，99 年度雜交 F2 後代植株農藝性狀調查結果，株高以臺東 1 號×S-14 之組合平均為 204.7cm 最高、成熟莢數以臺東 2 號×S-76 之組合平均為 376.3 個莢最高、單株粒重亦以臺東 2 號×S-76 之組合平均為 181.3g 最好、百粒重以臺東 3 號×S-95 之組合平均為 12.0g 最佳 (表 1)。

100 年度樹豆雜交 F1 後代植株，因 8 月 28 日中颱「天秤」颱風侵襲致臺東 1 號×S-14 之組合後代喪失，其餘植株農藝性狀調查結果，株高以臺東 2 號×S-76 之組合平均為 183.2cm 最高、成熟莢數亦以臺東 2 號×S-76 之組合平均為 472.2 個莢最高、單株粒重以 S-76×臺東 2 號之組合平均為 197.3g 最好、百粒重臺東 3 號×S-95 之組合平均為 11.2g 最佳 (表 2)。

表 1. 99 年度樹豆雜交 F2 後代植株農藝性狀調查

親本	株高 (cm)	株幅＋ (cm)		成熟莢數	未熟莢數	單株粒重 (g)	百粒重 (g)	籽粒顏色
臺東 1 號×S-14	204.7	146	142	195.5	32.9	105.9	11.8	白
S-14×臺東 1 號	186.7	161	142	318.6	6.5	159.5	10.5	白
臺東 2 號×S-76	181.0	159	140	376.3	6.2	181.3	10.7	茶褐
S-76×臺東 2 號	196.7	158	140	323.5	13.6	178.9	10.9	白
臺東 3 號×S-95	200.7	164	151	261.1	18.3	154.0	12.0	黑
S-95×臺東 3 號	201.7	165	149	330.0	4.3	176.7	11.3	白

採收日期：102 年 1 月 28 日
調查日期：102 年 3 月 5 日

表 2. 100 年度樹豆雜交 F1 後代植株農藝性狀調查

親本	株高 (cm)	株幅＋ (cm)		成熟莢數	未熟莢數	單株粒重 (g)	百粒重 (g)	籽粒顏色
臺東 1 號×S-14	－*	－	－	－	－	－	－	－
S-14×臺東 1 號	－	－	－	－	－	－	－	－
臺東 2 號×S-76	183.2	162	155	472.2	51.1	172.3	9.0	茶褐
S-76×臺東 2 號	177.8	143	136	367.5	28.6	197.3	10.9	白
臺東 3 號×S-95	179.9	165	153	360.7	25.9	164.2	11.2	黑
S-95×臺東 3 號	177.8	143	136	367.5	28.6	197.1	10.9	白

採收日期：102 年 1 月 7 日
調查日期：102 年 3 月 5 日
* 8 月 28 日中颱「天秤」颱風侵襲造成雜交後代植株全枯死。

101 年進行樹豆品種 (系) 雜交選育工作，由本場選育之臺東 1 號 (白色)、TS-221(紅色)、臺東 2 號 (茶褐色)、臺東 3 號(黑色)及 TS-303(紅色)等 5 品種 (系) 為親本，作為雜交的材料。以臺東 1 號×TS-221、臺東 2 號×TS-221 及臺東 3 號×TS303 等三個組合進行正、反雜交，共計 6 個組合。獲得雜交後代 F1 種子數為 1,625 個種子、成功率 23.6% (表 3)。記錄雜交期間日溫變化，日溫最低 15℃、日溫最高 28℃、日溫平均 20.5℃，詳如圖 1。進行人工雜交成功三天後略可看見幼莢，1 周後幼莢即可成形，幼莢生長 6-7 周後即可成熟，莢果授粉後生長情形如圖 2。探討樹豆授粉成功率原因可歸納為三大原因：一、氣候環境：包括溫度、濕度、日照、晴天與陰天。二、材料因素：包括取樣花苞成熟度、花粉活力、自然淘汰、蟲害影響。三、人為因素：去雄時人為操作不當而傷及花器、授粉量不足等。

期望於雜交後代能選育出籽實產量高、成熟期一致且產量、品質皆穩定之樹豆雜交品種，並適合於臺灣中、低海拔地區種植，以增加農民收益，進而促進臺東樹豆產業發展。

表 3.101 年度樹豆雜交授粉成功率及 F1 種子數

親本	授粉花朵數	莢果數	成功率(%)	種籽數
臺東 1 號×TS221	273	42	15.4	196
TS221 ×臺東 1 號	278	45	16.2	202
臺東 2 號×TS221	284	90	31.7	298
TS221 ×臺東 2 號	281	60	21.4	260
臺東 3 號×TS303	295	99	33.6	421
TS303×臺東 3 號	261	58	22.2	248
總　　計	1,672	394	23.6	1,625

授粉期間：自 101 年 11 月 28 日至 102 年 1 月 11 日止
調查日期：102 年 2 月 26 日




圖 1. 樹豆雜交授粉期間每日調查溫度變化圖 (日溫最低：15℃，日溫最高：28℃，日溫平均：20.5℃)




圖 2. 雜交授粉成功後莢果生長情形 (A：1 周、B：2 周、C：3 周 D：4 周、E：5 周及 F：6 周後)

引用文獻

Ae N. Arihara J, Okada K, Yoshihara T, Johansen C. 1990. Phosphorus uptake by pigeonpea and its role in cropping systems of the Indian subcontinent. Science. 248:477-480.

Amalraj, T., Ignacimuthu, S. 1998. Evaluation of the hypoglycaemic effect of *Cajanus cajan* (seeds) in mice. Indian J. Exp. Biol. 36: 1032-1033.

Grover, J. K., Yadav, S., Vats, V. J. 2002. Medicinal plants of India with anti-diabetic potential. J. Ethnopharmacol. 81:81-100.

Nan Wu, Kuang Fu, Yu-Jie Fu, Yuan-Gang Zu, Fang-Rong Chang, Yung-Husan Chen, Xiao-Lei Liu, Yu Kong, Wei Liu and Cheng-Bo Gu. 2009. Antioxidant activities of extracts and main components of pigeonpea (*Cajanus cajan* (L.)Millsp.) Leavs. Molecules. 14:1032-1043; dio:10.3390/molecules 14031032.

Saxena, K. B. 2008. Genetic improvement of pigeonpea-A Review. Tropical Plant Biol.1:159-178.

Sharma D., and J. M. Green. 1980. pigeonpea. In: Hybridization of crop plants. Amer Soc Agron Crop Sci. Society of America, Wisconsin, pp471-481.

Tang, Y., Wang, B., Zhou, X. J. 1999. Effect of external application of herbal cajani

Preparation on the fibronection content during healing process of open wound. J.

Guangzhou U. Tradit. Chin. Med.16: 302-304.

Wallis E. S., Byth D. E., Saxena K. B. 1981. Flowering responses of thirty-seven early maturing lines of pigeonpea. In: International workshop on pigeonpeas, vol.2: 15-19..

Hybridization and Breeding of Pigeon pea

Chen-I Chen

Taitung DARES, COA, Executive Yuan

Abstract

Pigeon pea is planted and cultivated generally in the areas of Taitung aboriginal tribal areas which is a kind of traditional indigenous crops, and also is regarded as health food. Presently, since there is a maturity inconsistent and low-yielding phenomenon on the domestic local strains which would cause the troubles in harvesting and management. In order to solve the issues above, intend to adopt the methods of artificial hybridization, emasculation and pollination during the period of pigeon pea flowering. Afterwards, to select out the new species which is high production and is consistent in maturity so as to provide the new hybrid varieties for farmers to plant. The success rate of the hybrid seeds by the way of artificial hybridization is 23.6% in this field in 2012, and expecting this new varieties could offer the framers to plant in order to increase their income in the future.

Key words: Pigeon pea, Hybridization, Breeding, Emasculation, Pollination.